

Synthesis , Characterization, and Biological Efficacy on new mixed ligand complexes based from azo dye of 8-hydroxy quinoline as a primary ligand and imidazole as a secondary ligand with some of transition metal ions

Israa N.Witwit^{1*}, Zahraa Y. Motaweq², Husham M.Mubark³

¹ University of Kufa , Faculty of Science , Dept. of Chemistry ,Najaf , Iraq

² University of Kufa , Faculty of Science , Dept. of Biology ,Najaf , Iraq

³ Islamic University, Clinical Laboratory Investigation Techniques, Najaf , Iraq.

Abstract:

New series of mixed ligand complexes of Mn(II), Co(II), Ni(II), Cu(II), and Hg(II) ions were prepared in two general formula [M (L₁)(L₂)₂]Cl , and [M (L₁)₂(L₂)₂] for each ion with E-5-((4-nitro phenyldiazany)quinoline-8-ol (L₁) as primary ligand , and imidazole molecule (L₂) as a secondary ligand . Free ligands and their complexes characterized via MS , UV-Vis. , FTIR , ¹HNMR, Magnetic susceptibility, and Molar Conductivity . The results indicating the octahedral geometry for all compounds with [M(L₁)₂(L₂)₂] formula while the complexes which have general formula [M(L₁)(L₂)₂]Cl articulate tetrahedral geometry except [Cu(L₁)(L₂)₂]Cl which has square planer geometry . (L₁) ligand behaved as bidentate through nitrogen atom and oxygen atom of hydroxyl group in the basic medium whereas imidazole coordinated through nitrogen (3) as a neutral monodentate ligand. Antibacterial efficiency of compounds were tested against (*Enterococcus faecalis* , *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Klebsiella pneumoniae*) multi-drug resistant bacteria (MDR). All compounds appeared significant efficacy contra MDR bacteria, but ligand L₁ appeared maximum inhibition zone and anti-bacterial efficacy contra all MDR bacteria while the minimum inhibition zone was marked in ligand L₂ antibacterial efficacy. The complexes generally have high antibacterial activities in gram +ve bacteria contrasted to gram -ve bacteria, where Hg (II) ion complexes showed higher biological efficacy on all bacteria than other mixed ligand complexes of other ions. Ni (II) ion complexes showed lower antibacterial activity compare to other metal complexes.

Key words: Azo compounds , 8-hydroxyquinoline , imidazole , mixed ligand compounds , biological efficacy.

INTRODUCTION:

Mixed ligand complexes of transition metal ions have great interesting from the researchers who study their coordination behavior and exploiting their properties in a different fields especially the antibacterial activity [1-3] , 8-hydroxy quinoline is an important compound which has the ability to coordinate with a various ions as bidentate through nitrogen atom of quinoline ring and oxygen atom after deprotonation of hydroxyl group [4-7] to form a five member ring between this ligand with the central metal ion that helps to increase the stability of the complexes , as well as its known biological activity [8] , Imidazole molecule is a part of the installation of many biological systems such as histidine and many of its derivatives have been used as antibacterial, antifungal and anticancer agents also the important using to estimation of metal ions in drugs [9,10] . This molecule has an ability to coordinate alone with metal ions as monodentate ligand through (N₃) atom [11] .

Pathogenic bacteria have caused dangerous illnesses and a lot of mortality in numerous countries, chiefly in the developing nations. These factors commonly diffuse speedily, and the common contagion to them have been appointed to the immune-compromised individuals, gravid women, children, and elderly persons [12]. Along with the gradual resistance of bacteria to the existing antibiotics as a consequence of irregular antibiotic exhaustion in medicine, the health and general hygiene of people are strongly at risk, and, hence, to avoid this threat, recognition and using new anti-bacterial compounds are requested [13]. In the

lately years, empirical research have showed some of the imidazole derivatives, numerous features such as the antifungal, antiviral, anti-parasitic, and in-vitro inhibition of cancer cells have been showed the antibacterial effects of these compounds has confirmed their capability to inhibit pathogenic bacteria such as *E. faecalis*, *S. aureus*, and *P. aeruginosa* [14,15].

The purpose of the present study is attended of mixed ligand complexes of azo dye of 8-hydroxy quinoline as a primary ligand and imidazole as a secondary ligand with Mn(II), Co(II), Ni(II), Cu(II), while Hg(II) in a different ratios of the primary ligand and characterize them by spectrometric techniques and examine the biological activity against different species of MDR bacteria .

MATERIALS AND METHODS

Chemicals and Instruments :

All chemical compounds used in this study were supplied by Merck , and B.D.H companies chemical company with 99% of purity , Electronic spectra measured by (UV-Vis)T80 , PG instruments Ltd. UK , while IR spectra recorded by (FTIR)-Platinum –ATR Bruker using KBr disk (400-4000)cm⁻¹ , Molar conductivity performed on 720(WTW) , Mass spectra was carried out by AB SCIEX 3200 QTRAP Mass analyzer , The element analysis was measured by Costech ECS Elemental 4010 , while magnetic measurements of complexes carried out by Balance Magnetic Susceptibility Model –M.S.B Auto , and ¹HMR in DMSO-d₆ solvent recorded by Bruker Avance-111 300 MHz NMR Spectrometer .

Preparation of (L₁) ligand

4-nitro aniline (1.38 gm, 10 mmole) dissolved in a mixture of (3 ml of HCl, and 15 ml of distilled water) in an ice bath with temperature below (5)^oC, then diazotized by adding a mixture of (0.70 gm NaNO₂, and 10 ml of cold distilled water) drop by drop with continuous stirring. The resulting diazonium salt was coupled with an alcoholic (0.44 gm NaOH, and 15 ml of ethanol) solution of (1.45 gm, 10 mmole) 8-hydroxy quinoline. After neutralization with diluted HCl solution the ligand was separated by filtration then dried in air. The yellow precipitate was produced with 79% of yield as shown in **scheme (1)**.

Preparation of complexes (general method)

The solid complexes were prepared by mixing (10) mmole of the primary ligand (L₁) for [M(L₁)(L₂)₂]Cl formula, and (20)mmole for [M(L₁)(L₂)₂] formula with (20) mmole of imidazole (L₂), and (10) mmole of metal chloride in (25) ml of absolute ethanol. The drops of (1)N of NaOH added to this mixture reflux the mixture for (2) hours until the precipitation was formed then filtered, dried in air, and recrystallized from ethanol. The schemes (2) and (3) show the general reaction to prepare the complexes, and table (1) illustrate some of physicochemical properties of ligands and its complexes.

Biological Activity

Bacterial Isolates:

The following multi drug resistance (MDR) pathogenic bacterial isolates: two gram +ve bacteria (*Staphylococcus aureus* and *Enterococcus faecalis*) while four gram -ve bacteria (*Proteus mirabilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*) were isolated from different clinical specimens like sputum, stool, CSF, wound, blood and urine. The isolates were diagnosed using a range of morphological as well as biochemical technicalities [16], and then finally confirmed by using Vitek-2 compact system GP and GN card automated bacterial identification instrument. All bacterial isolates were stocked on brain heart infusion broth with (15%) glycerol at (-20 °C). The isolates were sub-cultured on brain heart infusion agar and incubated at 37 °C for 24h before use.

Preparation of ligands and Complexes solutions:

The following concentrations were used in the antibacterial test:

1- Ligand concentration: 0.01 g of powder each ligand (L₁ and L₂) were dissolved in 1ml of DMSO to give concentration 10 mg/ml.

2- Mixed ligand complexes of ions: 0.01 g of powder of each complexes of Mn(II), Co(II), Ni(II), Cu(II) and Hg(II) for [M(L₁)(L₂)₂]Cl, and [M(L₁)₂(L₂)₂] formulas were dissolved in 1ml of DMSO to give 10mg/ml concentration.

Antibacterial activity experimental

Bacterial suspensions were prepared as explained by [17]. Agar well diffusion method used to determine the antibacterial activity of ligands and mixed ligand complexes of ions against bacterial isolates [18]. Brain heart infusion broth (BHIB) was used for the elaboration of MDR bacterial cultures. Muller Hinton agar (MHA) was used to determine the activity of ligands and mixed ligand

complexes of ions against bacterial. These media were attended similarity to the industrial's prescript.

Agar well diffusion assay

Bacterial isolates suspensions were attended to resemble 0.5 McFarland standards. Utilizing the micropipette, 100 μl of bacterial suspensions BHIB was diffuse up the superficies of MHA plate. This procedure was utilized for all experimental MDR bacteria. Utilizing an antiseptic cork borer, punctures were perforation in all of the culture plates. One of the punctures was perforation in the center of the plate where 100 μl of Gentamicin was inserted as positive control; 100μl of DMSO was inserted as a negative control in the other hole; 100μl of each ligands and mixed ligand complexes of ions were alone placed in the remaining holes [five holes to the [M(L₁)(L₂)₂]Cl complexes and five holes to other ratio [M(L₁)₂(L₂)₂] complexes. The culture plates were then incubated at 37°C for 24 h. The clear zone of inhibition around holes was calculated in mm. The tests were carried out in triplicate [19].


RESULTS AND DISCUSSION:

Mass spectra show a peak at m/e (294.3) represents the molecular ion peak of the free ligand (L₁), the primary fragmentation of (L₁) take place in two paths the first one by loose (N₂) of azo group at m/e(266.4) while the second path was started by loose (-OH) group at m/e (277) than loose (-N₂) from this fragment at m/e (249) as shown in scheme (4) and figure (1), Mass fragmentation of [Co(L₁)(L₂)₂]Cl complex that started by loose (-N₂) from (L₁), and [Co(L₁)₂(L₂)₂] complexes which fragmenting by loose two imidazole molecules respectively are being studied. The appearance of molecular peaks of these compounds gave agreement confirms of molecular formulas, as shown in figures (2, 3) and schemes (5, 6). ¹HNMR spectrum of (L₁) in DMSO-d₆ solvent show singlet at (8.86) ppm due to the proton of hydroxyl group of Quinoline moiety [20] which disappeared in [Hg(L₁)₂(L₂)₂] complex spectrum that indicate the coordination process through hydroxyl group after losing its proton, the fourth protons of aromatic aryl ring [21-24] obvious as a doublet signals in (6.62) and (6.65) ppm, while the protons of Quinoline appears at (7.10-8.3) ppm [25], The signal of (N1) proton of imidazole ring [26] was also observed at (12.25) ppm in the [Hg(L₁)₂(L₂)₂] complex spectrum, as shown in figures (4, 5).


IR spectra of (L₁) ligand show vibration frequencies of ν (C-O) for phenolic group [27, 28] in (1224) cm⁻¹ which proceed to higher frequencies in the complexes, as well the frequencies of this bond in plane and out of plane in (671) cm⁻¹ and (840)cm⁻¹ respectively, were priced to lower values in the complexes compared to free ligand, The ν (O-H) of phenolic group [29] that appear in (3473) cm⁻¹ minefield in the complexes, These above results indicated the coordination through oxygen atom of (-OH) group after it's deprotonation. The vibrational frequencies of ν (C=N) of Quinoline ring in (L₁), and ν (C=N) of imine group in (L₂) which appear in [30,31] (1506) cm⁻¹ and (1541) cm⁻¹ consecutively where observed to be shifted towards lower frequencies in the complexes If compared to the free

ligands suggesting that the coordination through nitrogen atoms of quinoline and imidazole rings of these ligands while there was no significant changes of $\nu(\text{N}=\text{N})$ in


complexes spectra which appear in (1444 cm^{-1}) [32] in (L1), as demonstrated in table (2).



Scheme(1): Preparation of (L₁)



Scheme (2) : Synthesis of $[\text{M}(\text{L}_1)(\text{L}_2)_2]\text{Cl}$ Complexes



Scheme (3) : Synthesis of $[\text{M}(\text{L}_1)_2(\text{L}_2)_2]$ Complexes




Figure 1: Mass spectrum of (*L*₁) ligand






Figure (2) : Mass spectrum of $[Co (L_1) (L_2)_2]Cl$ complex







Figure 3: Mass spectrum of Co (L₁)₂(L₂)₂ complex



Scheme (6) : Fragmentation of [Co (L₁)₂ (L₂)₂] Complex




Figure 4: ¹HMR spectra of (L₁) ligand




Figure 5 :¹HMR spectra of [Hg(L₁)₂(L₂)₂] complex




Figure 6: Uv-Vis spectra of (L₁)




Figure 8: Uv-Vis spectra of [Ni (L₁)(L₂)₂]Cl complex




Figure 7 :Uv-Vis spectra of (L₂)






Figure 9: Uv-Vis spectra of [Ni (L₁)₂(L₂)₂] complex



M= Mn(II) , Co(II), Ni(II) , Cu(II) , Hg(II)

Figure 10 :Suggested structure of [M(L₁)₂(L₂)₂] complexes




Figure 11: Suggested structure of $[M(L_1)(L_2)_2]Cl$ complexes




Figure 12: Comparing of antibacterial activity (inhibition zone, mm) of two ligands against pathogenic bacteria.




Figure 13: Antimicrobial activity activity (inhibition zone, mm) of mixed ligand complexes of Mn(II), Co(II), Ni(II) , Cu(II) , and Hg(II) ions against *Escherichia coli*.

Table(1) : Some of physicochemical properties of ligands and their complexes

Empirical Formula	Mwt	Yield(%)	Elemental Analysis Calcl. (Found)				m.p (°C)
			C%	H%	N%	M%	
C ₁₅ H ₁₀ N ₄ O ₃ (L ₁)	294.20	79	61.22 (61.20)	3.43 (3.41)	19.04 (19.06)	-----	124 – 126
C ₃ H ₄ N ₂ (L ₂)	68.07	-	52.93 (52.91)	5.92 (5.90)	41.15 (41.14)	-----	90 - 92
MnC ₂₁ H ₁₇ N ₈ O ₃ Cl	519.80	70	48.52 (48.68)	3.30 (3.32)	21.56 (21.52)	10.57 (10.59)	134-136
CoC ₂₁ H ₁₇ N ₈ O ₃ Cl	523.80	68	48.15 (48.20)	3.27 (3.24)	21.39 (21.18)	11.25 (11.28)	139-141
NiC ₂₁ H ₁₇ N ₈ O ₃ Cl	523.56	73	48.18 (48.20)	3.27 (3.26)	21.40 (21.42)	11.21 (11.16)	146-148
CuC ₂₁ H ₁₇ N ₈ O ₃ Cl	528.41	80	47.73 (47.58)	3.24 (3.26)	21.21 (21.08)	12.03 (12.07)	153-155
HgC ₂₁ H ₁₇ N ₈ O ₃ Cl	665.45	71	37.90 (37.94)	2.57 (2.60)	16.84 (16.85)	30.14 -----	166-169
MnC ₃₆ H ₂₆ N ₁₂ O ₆	777.61	75	55.60 (55.64)	3.37 (3.36)	21.62 (21.63)	7.07 (6.95)	189-191
CoC ₃₆ H ₂₆ N ₁₂ O ₆	781.60	81	55.32 (55.35)	3.35 (3.32)	21.50 (21.46)	7.49 (7.50)	216-219
NiC ₃₆ H ₂₆ N ₁₂ O ₆	781.36	76	55.34 (55.32)	3.35 (3.37)	21.51 (21.50)	7.51 (7.53)	221-225
CuC ₃₆ H ₂₆ N ₁₂ O ₆	786.21	70	55.00 (54.86)	3.33 (3.34)	21.38 (21.36)	8.08 (7.97)	226-228
HgC ₃₆ H ₂₆ N ₁₂ O ₆	923.26	77	46.83 (46.81)	2.84 (2.87)	18.21 (18.17)	21.73 -----	232-235

Table(2) : IR vibrations of (L₁) and (L₂) ligands and their metal ion complexes

Compound	ν (C-O) in plane	ν (C-O) out of plane	ν (C-O) phenolic	ν (C=N) imidazole	ν (C=N) quinoline	ν (M-N)	ν (M-O)
(L ₁)	671 w	840 w	1224 w	-	1506 m	-	-
(L ₂)	-	-	-	1541 m	-	-	-
[Mn (L ₁)(L ₂) ₂]Cl	659 w	836 m	1271 m	1546 m	1523 m	511 w	418 m
[Mn(L ₁) ₂ (L ₂) ₂]	657 w	823 w	1232 m	1464 m	1462 m	534 w	420 w
[Co (L ₁)(L ₂) ₂]Cl	634 w	821 w	1236 m	1550 m	1498 m	553 w	412 w
[Co(L ₁) ₂ (L ₂) ₂]	667 w	823 w	1228 w	1579 m	1496 m	504 w	414 w
[Ni (L ₁)(L ₂) ₂]Cl	661 w	842 m	1228 w	1548 m	1529 m	514 w	432 w
[Ni(L ₁) ₂ (L ₂) ₂]	665 w	821 w	1232 w	1570 m	1462 m	505 w	428 w
[Cu(L ₁)(L ₂) ₂]Cl	648 w	832 w	1234 w	1548 m	1467 m	518 w	414 w
[Cu(L ₁) ₂ (L ₂) ₂]	663 w	835 w	1230 w	1577 m	1470 m	520 w	432 w
[Hg (L ₁)(L ₂) ₂]Cl	651 w	842 m	1227 w	1552 m	1488 m	516 w	436 w
[Hg(L ₁) ₂ (L ₂) ₂]	663 w	826 w	1232 w	1564 m	1458 m	534 w	410 w

Molar Conductivity was carried out in both of (DMF) and (DMSO) solvents at (25)°C and (10⁻³)M for the prepared complexes , The resulting values for [M(L₁)(L₂)₂]Cl complexes were between (74.6-82.7) S.cm².mole⁻¹ in (DMF) solvent , and (72.8 – 81.4) S.cm².mole⁻¹ in (DMSO) indicating (1:1) electrolyte type , The white precipitate formed when a drops of (0.1) M of AgNO₃ was added to the complexes which confirms existence of chlorine out of the coordination sphere [33], while [M(L₁)₂(L₂)₂] showed values ranging between (23.5-17.5) S.Cm².mole⁻¹ in (DMF) , and (15.8-20.4) S.Cm².mole⁻¹ in (DMSO) confirmed the non-ionic character of these complexes [34] . Magnetic susbtibility measurements indicates that complexes formula [M(L₁)(L₂)₂]Cl have tetrahedral

geometry bating Cu(II) complex which has square planar while that all of [M (L₁)₂(L₂)₂] complexes were possessive the octahedral [35] .

Uv-Vis spectra of the complexes and free ligands measured in DMF solvent at room temperature , (L₁) resulted in two bands in 389 nm(25706 cm⁻¹) , and 256 nm (39062 cm⁻¹) due to the charge transfer and $\pi - \pi^*$ transitions while (L₂) showed two bands 287 nm (34843 cm⁻¹), and 226 nm(44247 cm⁻¹) consequent to n- π^* and $\pi - \pi^*$ transitions , All of these bands showed bathochromic shift in the complexes comparing to the ligands indicating the coordination process , while [Cu(L₁)(L₂)₂]Cl complex show single band at 616 nm (16233 cm⁻¹) due to ²B_{1g} ²A_{1g} transition of square planar geometry [34,36], The data of

transitions is summarized in table (3), and figures (6 - 9) represent the electronic spectra of the free ligands and Ni(II) complexes

The capacity of antimicrobial efficiency of two ligand (L₁,L₂) and five mixed ligand complexes of Mn(II), Co(II), Ni(II), Cu(II), and Hg(II) in two formulas [M(L₁)(L₂)₂Cl] and [M(L₁)₂(L₂)₂] for each ion against six multidrug resistance bacteria (two G +ve bacteria (*E. faecalis* and *S. aureus*) and four G –ve bacteria (*P. mirabilis*, *E. coli*, *P. aeruginosa* and *K. pneumoniae*) were assessed by the existence or the absence of inhibition zone. The results of antibacterial efficiency of the ligands and mixed ligand complexes of ions are listed in table (4), table (5) and figure (13). The results indicate that L₁ ligand had better antibacterial activities on all bacterial isolated than L₂ ligand and other mixed ligand complexes of ions. The results also indicated that L₂ ligand have no influence on G +ve MDR bacterial isolates compared to the influence on G –ve MDR bacterial isolates .

As for the effectiveness of the metal complexes, they generally have a high antibacterial activities on G +ve bacteria compared to G-ve bacteria, where Hg (II) ion complexes had higher biological efficacy on all bacteria

than other metal complexes. Ni (II) ion complexes showed lower antibacterial activity compare to other metal complexes (table 5).

The complexes [M(L₁)₂(L₂)₂] of Cu(II), Mn(II), Ni(II) and Co(II) ions given higher antibacterial activity against all bacteria compare to [M(L₁)₁(L₂)₂Cl] except [Hg(L₁)₂(L₂)₂] of complex showed lower antibacterial activity compare to [Hg(L₁)(L₂)₂Cl] (figure 13).

The enhancement of antimicrobial activity of the ligand (L₁) may be due to chelation of transitional metals with it. Complicated decrease the polarity of metal ion by coordinating with ligands and rises the lipophilicity of the metals [37]. Thus it easiness the new synthesized complex to permeate the lipid cell membrane of microorganisms and prevent their growth.

This variation in efficacy and inhibition vigor of(L₂) ligand is compatible to the existence of chlorine and phenyl compounds [38]. Beside the reduction of influences on G +ve bacteria, experiments demonstrated the potency of methyl nitro imidazole to inhibit the growth of the G-ve bacteria such as *Proteus merablis*, *P. auroginosa*, *K. pneumoniae* and *E. coli*, This ligand could liberate free radicals that damage bacteria and murder them [39] .

Table(3) : Molar Conductivity, Magnetic Susbtibility , and Electronic Transitions of ligands and their complexes .

Compound	Molar Conductivity S.cm ² .mole ⁻¹		μ.eff. (B.M.)	λ max (nm)	Transitions	Geometry
	DMF	DMSO				
(L ₁)	-----	-----	-----	256 389	π-π* C.T	-----
(L ₂)	-----	-----	-----	226 278	π-π* n-π*	-----
[Mn (L ₁)(L ₂) ₂]Cl	78.8	75.6	5.93	238 262 422	π-π* π-π* MLCT	Tetrahedral
[Mn(L ₁) ₂ (L ₂) ₂]	20.1	18.3	5.72	232 259 418	π-π* π-π* MLCT	Octahedral
[Co (L ₁)(L ₂) ₂]Cl	76.4	74.7	3.87	234 264 432	π-π* π-π* MLCT	Tetrahedral
[Co(L ₁) ₂ (L ₂) ₂]	18.6	16.5	4.72	230 258 429	π-π* π-π* MLCT	Octahedral
[Ni (L ₁)(L ₂) ₂]Cl	76.2	74.5	3.53	236 267 435	π-π* π-π* MLCT	Tetrahedral
[Ni(L ₁) ₂ (L ₂) ₂]	18.4	16.4	2.85	232 260 432	π-π* π-π* MLCT	Octahedral
[Cu(L ₁)(L ₂) ₂]Cl	74.6	72.8	1.81	267 456	π-π* MLCT	Square planar
[Cu(L ₁) ₂ (L ₂) ₂]	17.2	15.8	1.71	264 448	π-π* MLCT	Distorted octahedral
[Hg (L ₁)(L ₂) ₂]Cl	82.7	81.4	-----	271 256 478	π-π* π-π* MLCT	Tetrahedral
[Hg(L ₁) ₂ (L ₂) ₂]	23.5	20.4	-----	268 480	π-π* MLCT	Octahedral

Table (4): Antibacterial activity (inhibition zone, mm) of ligands against pathogenic bacteria

Type of bacteria	L1	L2
<i>Staphylococcus aureus</i>	35	0
<i>Enterococcus faecalis</i>	45	0
<i>Escherichia coli</i>	50	10
<i>Klebsiella pneumoniae</i>	45	10
<i>Proteus mirabilis</i>	56	15
<i>Pseudomonas aeruginosa</i>	35	8

Table (5): Antibacterial activity (inhibition zone, mm) of mixed ligand complexes of Mn(II), Co(II), Ni(II), Cu(II), and Hg(II) ions against pathogenic bacteria

Compound	<i>S. aureus</i>	<i>E. faecalis</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>P. mirabilis</i>	<i>P. aeruginosa</i>
Mn (L ₁)(L ₂) ₂ Cl	25	25	25	7	0	16
Mn(L ₁) ₂ (L ₂) ₂	30	27	29	15	14	18
Co (L ₁)(L ₂) ₂ Cl	15	17	0	2	0	0
Co(L ₁) ₂ (L ₂) ₂	24	19	13	22	20	28
Ni (L ₁)(L ₂) ₂ Cl	20	0	0	0	0	0
Ni(L ₁) ₂ (L ₂) ₂	23	0	11	0	11	0
Cu (L ₁)(L ₂) ₂ Cl	40	0	19	7	24	5
Cu(L ₁) ₂ (L ₂) ₂	40	13	27	12	28	3
Hg (L ₁)(L ₂) ₂ Cl	34	26	30	25	25	24
Hg(L ₁) ₂ (L ₂) ₂	28	24	27	22	14	21

CONCLUSIONS

Series of new mixed ligand complexes were prepared with (E)-5-nitrophenyl)diazanylquinolin-8-ol as a primary ligand (L₁) and imidazole as a secondary ligand (L₂) with Mn(II), Co(II), Ni(II), Cu(II), and Hg(II), in two different ratios of (L₁) as two formulas [M(L₁)(L₂)]Cl which have tetrahedral geometry except [Cu(L₁)(L₂)]Cl has square planer and [M(L₁)₂(L₂)₂] were octahedral geometry.

The ligand (L₁) has maximum inhibition zone and antibacterial efficacy against all tested MDR bacteria while the minimum inhibition zone was determined ligand L₂ antibacterial efficacy. Retained The complexes generally appeared high antibacterial activities in G +ve bacteria compared to G -ve bacteria, where Hg (II) ion complexes has higher biological efficacy on all bacteria than other mixed ligand complexes of other ions. Ni (II) ion complexes has lower antibacterial activity compare to other metal complexes.

REFERENCES :

[1] Sunil S. Patil, Ganesh A. Thakur, Manzoor M. Shaikh, Synthesis, Characterization, and Antibacterial Studies of Mixed Ligand Dioxouranium Complexes with 8-Hydroxyquinoline and Some Amino Acids, *International Scholarly Research Network*, 2011, 168539, 1-6.

[2] Taghreed H AL-Noor, Amer J Jarad, Abaas Obaid, Synthesis, Characterization and antibacterial activity of mixed ligands complexes of some metal ions with 2-aminophenol and tributylphosphine, *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2017, 8(3), 132-139.

[3] F. K. Camellia, Abdul Kader, Md. Ashraf Alam, Md. Kudrat-Zahan and M. S. Islam, Synthesis, Characterization and Antimicrobial Investigation of Three New Mo (VI) Mixed Ligand Complexes, 2018; 4(4) : pp.1-5.

[4] R. T. Vashi, S. B. Patel, H. K. Kadiya, Synthesis, Characterization and Antimicrobial Activity of Metal Chelates of 2-[(8-hydroxyquinolinyl)-5-aminomethyl]-3-(4-Bromophenyl)-3(H)-quinazolin-4-one, *Der Pharma Chemica*, 2012, 4(4):1506-1511

[5] R.B. Dixit, J.A. Patel, SPECTROPHOTOMETRIC DETERMINATION OF PARACETAMOL DRUG USING 8-HYDROXYQUINOLINE, *International Journal of Pharmaceutical Sciences and Research*, 2014; 5(6): 2393-2397.

[6] Ibrahim M.A. Awad, Aref A.M. Aly, Aly A. Abdel Hafez, Krm. Hassan, Transition Metal Complexes of Azo-8-Hydroxyquinoline Derivatives as Antimicrobial Agents, *Journal of the Chinese Chemical Society*, 1989; 36: 107-114.

[7] Nadia A. El - Wakiel, Hala F. Rizk, Seham A. Ibrahim, Synthesis and characterization of metal complexes of azo dye based on 5-nitro-8-hydroxyquinoline and their applications in dyeing polyester fabrics, *Appl Organometal Chem*, 2017;31(10):1-7.

[8] Mohamed A. Abdelgawad, Ashraf M. Mohamed, Arafa Musa, Ehab M. Mostafa, Hassan M. Awad, Synthesis, Chromatographic Separation and Antimicrobial Evolution of New Azoquinoline-8-ol, *J. Pharm. Sci. & Res.*, 2018; 10(6) : 1314-1318.

[9] Sanchita Baroniya, Zaihra Anwer, Pramod K. Sharma, Rupesh Dudhe, Nitin Kumar, Recent advancement in imidazole as anti cancer agents: A review, *Der Pharmacia Sinica*, 2010, 1 (3): 172-182.

[10] Arshed A. Ali Shihad, Khamis A. Abedalrazaq, Faiq F. Karam, Haider M. Hessoon, Khalid J. Al-Adilee, Ayad F. Alkaim, New Analytical Method for Estimation of ferrous in Ferrous Sulfate drug By Preparation and Using 2-(E-(1H-benzo(d)imidazol-2-yl) diazenyl-5-(E-4-dimethyl amino benzaliden amino) phenol as a reagent, *J. Pharm. Sci. & Res.*, 2018, 10: 92179-2182

[11] J. Reedijk, Pyrazoles and Imidazoles as Ligands. II. Coordination Compounds of N-methyl Imidazole with Metal Perchlorates and Tetrafluoroborates, *Inorganica Chimica Acta*, 1969; 3(4) :517-522.

[12] Giesker K, Hensel M. Bacterial Vaccines, *Ref Mod Bio Sci*, 2014; doi:10.1016/B978-0-12-801238-3.00141-0.

[13] AL-Ramahi, A.A.; Darweesh, M. A., Ahmad A. M (2017).: The Antibacterial of Essential Fatty Acid Semicarbazide Extracted from Flaxseed Oil Against Some Nosocomial Infection Bacteria in Iraq. *IJCPR*; 8(1): 31-39.

[14] Brahmayya M, Venkateswararao B, Krishnarao D, Durgarao S, Viplava Prasad U, Damodharam T, Synthesis and fungicidal activity of novel 5-aryl-4methyl-3yl (imidazolidin-1yl methyl, 2-ylidene nitro imine) isoxazoles., *J Pharm Res*, 2013;7: 516-519.

[15] Salhi L, Bouzroua-Aichouche S, Benmalek Y, Benmalek B, Poulain-Martini S, Cacciuto B, An efficient conversion of maleimide derivatives to 2-thioxo imidazolidinones. *Org Commun*, 2013;6: 87-94.

- [16] Macfaddin, J.F. Biochemical tests for identification of medical Bacteria. 3rd-ed, willium and Wilkins, U. S. A., 2009.
- [17] Ramalivhana JN, Obi CL, Samie A, Iweriebor BC, Uaboi-Egbenni P, Idighe J.E. and Momba M. N. B., Antibacterial activity of honey and medicinal plant extracts against Gram negative microorganisms, *African Journal of Biotechnology*, 2014; 13(4): 616-625.
- [18] Murray, PR. ; Baron EJ , Pfaller MA, Tenover FC , Tenover HR., Manual of clinical microbiology, 6th Ed. ASM press , Washington DC, 1995; 15-18.
- [19] Olurinola, P.F. A laboratory manual of pharmaceutical microbiology. Idu, Abuja, Nigeria, 1996; 69-105.
- [20] Chinnagiri T. Keerthi Kumar, Jathi Keshavayya, Tantry N. Rajesh, Sanehalli K. Peethambar, Angadi R. Shoukat Ali, Synthesis, Characterization, and Biological Activity of 5-Phenyl-1,3,4-thiadiazole-2-amine Incorporated Azo Dye Derivatives , *Organic Chemistry International*, 2013; ID 370626, 1-7.
- [21] M. El Faydy, T. Djassinra, S. Haida, M. Rbaal, K. Ounine, A. Kribii, B. Lakhriissi, Synthesis and investigation of antibacterial and antioxidants properties of some new 5- substituted-8-hydroxyquinoline derivatives, *Journal of Materials and Environmental Sciences*, 2017 ; 8(11): 3855-3863
- [22] Won Young Kang , Jong S. Park, Preparation of Polymeric Metal Complex Containing Azo Dye Rotaxane , *Textile Coloration and Finishing* , 2011, 23(3): 163-168.
- [23] Hitendra D. Raj ,Yogesh S. Patel, Synthesis, characterization and antifungal activity of metal complexes of 8-hydroxyquinoline based azo dye, *Advances in Applied Science Research*, 2015, 6(2):119-123
- [24] K. B. Patel , G. J. Kharadi ,K. S. Nimavat, Synthesis and description of transition metal complexes and antimicrobial Studies, *Journal of Chemical and Pharmaceutical Research*, 2012, 4(5):2422-2428.
- [25] Hitendra D. Raj ,Yogesh S. Patel , Synthesis, characterization and antifungal activity of metal complexes of 8-hydroxyquinoline based azo dye , *Adv. Appl. Sci. Res.*, 2015; 6(2):119-123.
- [26] Mesut Gomleksiza , Cihan Alkanb and Belgin Erdemc, Synthesis Characterization and Antibacterial Activity of Imidazole Derivatives of 1,10-Phenanthroline and their Cu(II), Co(II) and Ni(II) Complexes, *S. Afr. J. Chem.*, 2013; 66:107–112.
- [27] Omar Hamad Shihab Al-Obaidi , Binuclear Cu(II) and Co(II) Complexes of Tridentate Heterocyclic Schiff Base Derived from Salicylaldehyde with 4-Aminoantipyrine , *Bioinorganic Chemistry and Applications*, 2012 ; 2012 : 1-6, doi:10.1155/2012/601879.
- [28] Masoumeh Orojloo, Fereshteh Nourian, Raziye Arabahmadi and Saeid Amani, Ni(II), Cu(II), AND Zn(II) COMPLEXES DERIVED FROM A NEW SCHIFF BASE 2-((Z)-(3-METHYLPYRIDIN-2-YLEIMINO)METHYL)PHENOL AND SYNTHESIS OF NANO SIZED METAL OXIDE PARTICLES FROM THESE COMPOUNDS, *Quim. Nova*, 2015; 38(9): 1187-1191, 2015.
- [29] Mosaad R. Mlahi, Elsayed M. Afsah, Amr Negm , Mohsen M. Mostafa, Synthesis of 8- hydroxyquinolium chloroacetate and synthesis of complexes derived from 8-hydroxyquinoline, and characterization, density functional theory and biological studies, *Appl. Organometal. Chem.*, 2015; 29:200–208.
- [30] Kuruba Siddappa , Nabiya Sultana Mayana, Synthesis, Spectroscopic Characterization, and Biological Evaluation Studies of 5-Bromo-3-((hydroxy-2-methylquinolin-7-yl)methylene) hydrazono)indolin-2-one and Its Metal (II) Complexes , *Bioinorganic Chemistry and Applications*, 2014; Article ID 483282: 11 pages , doi.org/10.1155/2014/483282.
- [31] John McGinley, Malachy McCann, Kaijie Ni, Theresa Tallon, Kevin Kavanagh, Michael Devereux , Xiaomei Ma, Vickie McKee , Imidazole Schiff base ligands: Synthesis, coordination complexes and biological activities, *Polyhedron* , 2013; 55: 169–178
- [32] Saad M Mahdi , Ali K Ismail, Preparation and Identification of new azo-schiff base ligand (NASAR) and its divalent transition metal Complexes , *J. Pharm. Sci. & Res.* ,2018; 10(9) , : 2175-2178.
- [33] Khalid J. Al-Adilee , Hussein A. K. Kyhoiesh, Preparation and Identification of Some Metal Complexes with New Heterocyclic Azo Dye Ligand 2-[2' - (1- Hydroxy -4- Chloro phenyl) azo]-Imidazole and their Spectral and Thermal Studies, *Journal of Molecular Structure* , 2017 ; DOI: 10.1016/j.molstruc.2017.01.054.
- [34] K. Hussain Reddy, M. Radhakrishna Reddy , K. Mohana Raju , Synthesis, characterization, electrochemistry and axial ligation properties of macrocyclic divalent metal complexes of acetylacetone buckled with different diamines , *Polyhedron* , 1997; 16(15) : 2673-2679.
- [35] C. Anitha, S. Sumathi, P. Tharmaraj, C. D. Sheela, Synthesis, Characterization, and Biological Activity of Some Transition Metal Complexes Derived from Novel Hydrazone Azo Schiff Base Ligand, *International Journal of Inorganic Chemistry* ,2011, Article ID 493942:8 pages , doi:10.1155/2011/493942
- [36] Sheikh J, Juneja H, Ingle V, Ali P, Hadda TB. Synthesis and in vitro biology of Co(II), Ni(II), Cu(II) and Zinc(II) complexes of functionalized betadiketone bearing energy buried potential antibacterial and antiviral O,O pharmacophore sites, *J Saudi Chem Soc.* 2013;17(3):269-76.
- [37] Franz A. Mautner, Roland C. Fischer, Mark Spell, Andres R. Acevedo, Diana H. Tran , Salah S. Massoud, Metal(II) Complexes of Compartmental Polynuclear Schiff Bases Containing Phenolate and Alkoxy Groups, *Crystals* ,2016; 6(91) , doi:10.3390.
- [38] Jones RN, Wilson ML, Weinstein MP, Stilwell MG, Mendes RE, Contemporary potencies of minocycline and tetracycline HCL tested against Gram-positive pathogens: SENTRY Program results using CLSI and EUCAST breakpoint criteria, *Diagn Microbiol Infect Dis* ,2013; 75: 402–405.
- [39] Shahid HA, Jahangir S, Yousuf S, Hanif M, Sherwan SK. Synthesis, crystal structure, structural characterization and in vitro antimicrobial activities of 1-methyl-4-nitro-1H-imidazole. *Arab J Chem* ,2014; doi:10.1016.