



Identification, Quantitative Determination & Antidepressant Activity of Chlorogenic Acid & Gallic Acid from *Morus alba* Leaves

MerajAli¹, AadityaSingh², Shiv Bhadra Singh³, Urmila Nishad⁴

Faculty of Pharmacy Gaya Prasad Institute of Human Excellence, Malihabad, Lucknow, Uttar Pradesh, India^{1,4}.

Faculty of Pharmacy Aryakul Group of Colleges, Lucknow, Uttar Pradesh, India²

Hygia Institute of Pharmacy, Lucknow, Uttar Pradesh, India³

Abstract

Objective: The present study is considered to estimate preliminary phytochemical components, physicochemical parameters, HPTLC chromatographic studies and antidepressant activity of chlorogenic acid & gallic acid present in extracts of *Morus alba* leaves.

Methods: Total phenol content, Total flavonoid content, Total tannin content, HPTLC fingerprinting analysis for compounds like chlorogenic acid & gallic acid responsible for antioxidant activity and antidepressant activity by TST & FST.

Result: Preliminary phytochemical studies show the presence of good phenol (21.75 ± 1.21) and flavonoid (14.83 ± 2.34) content. HPTLC fingerprinting by comparing extract values with standards available of chlorogenic acid & gallic acid also shows the presence of these components in the methanolic and ethanolic extracts of *Morus alba* leaves, which were further used to screen antidepressant activity and shows substantial antidepressant activity in the plant.

Conclusion: Our results recommend that *Morus Alba* leaves are may be demonstrated to be an acceptable natural antioxidant and antidepressant with various bioactive components used for the treatment of numerous other ailments.

INTRODUCTION

Natural plants are used as very good source of nutrition persistent food as well as source of various chemical constituents operative in curing various diseases which may demand as the biologically active constituents. At the present natural plants are very much in petition in the form of drugs because of their fewer side effects, they are considered the potential resources of various bioactive compounds and are also easily available from the natural sources. In the same context *Morus alba*, the Mulberry plant which is basically famous for sericulture, the fabrication of silk done through the silkworm and the leaves are also used to diminish the symptoms of diabetes in vernacular medicine as well as for improving cardio-metabolic risks, including antihyperglycaemic, antihyperlipidaemic, antiobesity, antihypertensive, antioxidative, anti-inflammatory, anti-atherosclerotic and cardioprotective effects¹ in Chinese medicine used to treat constipation, to tonify the blood, prematurely grey hair, cough, edema, to promote urination, fever, headache, dry & sore eyes² and so many more. So, the leaves is used further in this study to explore some more about the biological activity of leaves.

Mulberry plant belongs to genus *Morus* having 68 species which are unisex flowering plants belonging to family Moraceae of the Urticales subclass. The plant is a shrub or tree (20 to 30 feet high) often the size of a small apple tree, having leaves which are thin, glossy, and light green in color with 5 lobes or has one lobe, two lobes, three lobes, or no lobes at all. *Morus alba* L. is also known as White Mulberry and can be grown from seed as well as planted from large cuttings of root readily. Commonly, the plantation is upraised in a block foundation with arrangement of 6 feet x 6 feet, or 8 feet x 8 feet, as plant to plant and row to row spaces. The plants are generally trimmed once a year during the monsoon season (July –

August) to a height of 5–6 feet and allowed to grow with a maximum of 8–10 shoots at the top.³ The plant is widely distributed in India, China, Japan, North Africa, Arabia, South Europe, etc.

Morus alba L. leaves had been used as substantial source of medicine, drink, and functional foods in many countries. It is used in drinks as green tea with several other herbal drugs like tulsi and aswagandha because of its immune boosting antioxidants like Chlorogenic acid, rutin, isoquercitrin, and astragalin. Anticancerous alkaloids like 1-deoxynojirimycin, morroles B–F⁴, (2R,3R,4R)-2-hydroxymethyl-3,4-dihydropyridine-N-propionamide from the root bark and 4-O-R-D-galactopyranosyl-calystegine B2 and 3 β ,6 β -dihydroxynortropine from the fruits⁵, mulbaines A, B & C⁶. Eighteen important amino acids with calcium, potassium, sodium, magnesium, zinc, iron, copper, manganese, chromium, selenium, arsenic, vitamins and it's no caffeine property. Other chemical constituents present in leaves are coumarins, flavonoids, anthocyanins and polyphenols including quercetin 3-(malonylglucoside), rutin, isoquercitrin, cyaniding-3-rutinoside apigenin, luteolin, quercetin, morin, caffeic acid, gallic acid, umbelliferone, chlorogenic acid, and kaempferol.⁷ The plant extract rich in polyphenols used as a non-toxic natural healing agent, which also have high prospective applications as skin-whitening agents due to its potent tyrosinase inhibitor property.⁸

Chlorogenic acid is a phenolic compound which is an ester of caffeic acid. It is not stable thermally and is readily disintegrated to quinic acid and caffeic acid. Gallic acid also known as 3,4,5-trihydroxybenzoic acid is also a phenolic compound which usually bonded to form dimers such as ellagic acid. Both of these phenolics and their derivatives have received much consideration due to recent studies presenting different biological

properties of these classes of compounds. Both categories play very important role as antioxidants, for improving symptoms of diabetes as well as for direct and indirect prevention and cure of various other diseases. Hence both these compounds were chosen in this study to analyze and quantify by HPTLC fingerprinting, spectral analysis, elemental analysis to confirm the structure and to observe its antidepressant activity.




Fig 1. Structures of (a)-chlorogenic acid, (b)-Gallic acid

MATERIALS AND METHODS

Instruments & apparatus like digital balance of Citizen, UV chamber of Systronics, UV-Visible spectrophotometer of Systronics, Hot air oven of Science Tech, Heating mental of Science Tech and pH meter of Systronics were used at different steps as per requirements. Chemical used like such as petroleum ether, methanol, ethanol, Folin-Ciocalteus reagent, gallic acid, tannic acid, rutin, Na₂CO₃, aluminium chloride, NaNO₃, NaOH, were of analytical grade and purchased commercially from S D Fine, Mumbai. Glass wares like Soxhlet apparatus, conical flask, beaker, measuring cylinder, RBF, separating funnel, volumetric flask, test tube etc. were used of Borosilicate.

Crude Extract Preparation

Morus alba leaves were collected from local Lucknow and were authenticated by CSIR-NBRI. Fresh leaves were washed and dried through air at room temperature. After two weeks of air drying, leaves are crushed with the help of Mixture grinder. The powdered material used for extraction by Soxhlet apparatus using the solvents like petroleum ether, ethanol and methanol successively. During extraction process a temperature of 40- 60⁰ C was maintained for 6 hours. The concentrated product of extracted material was collected and stored in refrigerator for further experimental analysis.

Phytochemical screening

Preliminary phytochemical screening was done to investigate the phyto-constituents present in petroleum ether, ethanol and methanol extracts. Performed according to methods & procedures given in Practical Pharmacognosy book of C. K. Kokate.⁹ Results were tabulated in Table-1. The extract obtained by several extraction give positive test for alkaloids, carbohydrate, caumarin, phenolic compounds, tannins, proteins & amino acids etc.

Physicochemical studies

The physicochemical parameters like extract characteristics (consistency, color in day light, fluorescence analysis & % yield), moisture content, Chlorophyll content, ash value etc. in extracts were performed using methods reported in AOAC 1990¹⁰. Quantitative phytochemical analysis for total phenol content, total flavonoid content, total tannin content were also estimated and results were tabulated. Gallic acid, Rutin and tannic acid were taken as standards for estimation of total phenol, flavonoid and tannin content respectively.

HPTLC Chromatography Densitometry

Five working standard solutions from methanol were freshly prepared in concentration ranging from 0.1 – 0.5 mg/ml for calibration curve from stock solution of standards of 1 mg/mL. The working solution of standards (1mg/ml) and samples (10mg/ml) were freshly prepared in similar solvent.

The mobile phase used for development of HPTLC plates was toluene: ethyl acetate: formic acid in a ratio of 7: 2.5: 0.5 v/v for gallic acid and ethyl acetate: acetic acid: formic acid: water in a ratio of 10:1.1: 1.1: 2.3 v/v for chlorogenic acid. The analysis was carried out using Camag HPTLC system equipped with Linomat-V applicator and 100 µl syringe. The samples were spotted against the standards using microliter syringe over the pre-coated silica gel 60 F254 HPTLC plates, and development of the applied plate was carried out in pre-saturated Camag twin-trough chamber. Then the plates are dried and visualized in UV light of 254nm and 366nm wavelength and the results are interrelated in Tables 4-5 and Fig.2-3. The percentage of chlorogenic acid and gallic acid was calculated by using the formula.¹¹

$$\frac{(\text{sample area} \times \text{standard dilution} \times \text{purity}) \times 100}{(\text{standard area} \times \text{sample dilution} \times 100)}$$

Antidepressant Activity

Animals

Albino mice of either sex were acquired from the Central Animal House, Aryakul College of Pharmacy & Research (Reg.No- 1896/PO/Re/S/16/CPCSEA). Thirty swiss albino mice (25-35 gms body weight) of either sex were randomly selected and grouped into 5 groups (n=6). They were accustomed and housed in animal house with 12hr: 12hr light-dark cycle at 27±2^oc temperature and 45-55% RH. Food and water delivered ad libitum. The work was permitted by the Institutional Animal Ethical Committee (IAEC). Control animals were doped with distilled water.

Drugs like imipramine (10mg/kg), test drug *Morus alba* leaves extract (100 mg, 200mg, 400mg/kg) were dissolved in distilled water and doped orally once daily for 7 days (one week). On 8th day tests were reiterated.

Acute toxicity studies

The process for acute toxicity of ethanolic leaf extract of *Morus alba* was monitored as per the OECD guideline no.423 (Acute Toxic Class Method) (OECD, 2002). A dose of 800, 1000, 2000mg/kg bodyweight is overseen and animals were observed for 15 days.

Tail suspension test (TST): A mouse was drooped on a wire in an upside down position so that its nostrils just touch the water surface in a vessel. After initial enthusiastic movement, the mouse undertakes a motionless

posture and the period of motionless posture during five minutes reflection was noted. This test is consistent and a prompt screening method for antidepressants, including those involving the serotonergic system.¹²

Forced swimming test (FST): The rats were placed a cylinder (45x20cm) holding 38 cm water (25±2°C), so that the rat could not touch lowermost part of cylinder with its hind limb or tail or climb over the verge of the chamber. Two swim periods were accompanied, an initial fifteen minutes pre-test, followed by five minutes test 24 hour later. Drug was doped after pre-test. The period of motionlessness (remained floating in water without harassed and making only those movements necessary to keep its head above water) during five minutes test period was noted.¹³

RESULT & DISCUSSION

Table 1. Preliminary phytochemical screening of Pet ether, ethanolic & methanolic extract of *Morus alba* leaves

Constituents	Pet. Ether extract	Ethanolic extract	Methanolic extract
Alkaloids	-	++	+++
Carbohydrates		+	++
Caumarin	+	++	+++
Flavonoids	+	+++	++
Fixed oil	++	-	-
Glycosides	-	+	+
Gums and resins	-	-	-
Mucilages	-	+++	+++
Proteins & amino acid	-	++	+++
Saponins	-	-	-
Steroids	+	-	-
Tannins	-	+++	++

+++ High, ++ Moderate: + Slight: - Negative

Table 2. Extract characteristics

Types of solvent	Consistency	Colour in day light	Fluorescence analysis		% w/w
			Short UV	Long UV	
Ethanol	Semi-solid	Brownish black	Reddish black	Brownish black	5.82
Ethanol	Solid	Reddish brown	Reddish brown	Brown	6.32
Methanol	Solid	Brown	Greenish brown	Brown	7.14

Table 3. Physiochemical studies & quantitative phytochemical analysis

S.No.	Physiochemical Parameters	Values
1.	Moisture contents	3.1 ± 0.3%
2.	Ash value	12.1 ± 0.04% w/w
3.	Chlorophyll content	1.97 ± 0.03
4.	Total phenolics (mg gallic acid equivalent/g)*	21.75 ± 1.21
5.	Total flavonoid (mg rutin equivalent/g)*	14.83 ± 2.34
6.	Total tannin (mg tannic acid equivalent/g)*	7.21 ± 1.75

(*mean ± S.D, n=3)

Table 4.HPTLC- Phenols profile of methanolic extract of Morus alba leaves for chlorogenic acid

Peak	R _f	Height (AU)	Area (AU)	Assigned substances
1	0.01	612.1	14992.1	Unknown
2	0.04	603.5	8821.9	Unknown
3	0.06	515.2	26092.5	Chlorogenic acid
4	0.14	328.3	12914.6	Unknown
5	0.20	602.5	20777.9	Unknown
6	0.25	250.1	4616.3	Unknown
7	0.27	450.6	13821.7	Unknown
8	0.31	456.5	11207.6	Unknown
9	0.36	298.3	12070.3	Unknown
10	0.43	166.4	5633.1	Unknown
11	0.47	167.5	4345.4	Unknown
12	0.51	136.3	3452.4	Unknown
13	0.58	133.8	3561.1	Unknown
14	0.61	133.1	4753.1	Unknown
15	0.65	114.3	3271.5	Unknown
16	0.69	151.6	6863.0	Unknown
17	0.75	203.0	10216.0	Unknown
18	0.82	188.1	6897.9	Unknown
19	0.87	192.5	7083.6	Unknown

Table 5. HPTLC- Phenols profile of methanolic extract of Morus Alba leaves for gallic acid

Peak	R _f	Height (AU)	Area (AU)	Assigned substances
1	0.05	27.4	842.4	Unknown
2	0.47	18.7	196.6	Unknown
3	0.67	38.3	959.1	Gallic acid
4	0.71	72.3	1855.8	Unknown
5	0.77	167.7	6291.6	Unknown
6	0.83	192.5	10855.3	Unknown
7	0.93	14.4	346.6	Unknown




Fig. A.





Fig. B.

**Fig 2. HPTLC densitogram for methanolic extract with their respective standards:
A. Chlorogenic acid B. Gallic acid**



A. Chlorogenic acid B. Gallic acid
 Fig 3. 3D diagram of HPTLC densitograms

Results of Antidepressant activity:

Table 1: Effect of *Morus alba* leaves extract on immobility time in tail suspension test

Treatment	DOSE (mg/kg)	Immobility time (in sec)
Vehicle Control(6)	-	180.40±2.70
<i>M.alba</i> (6)	100	161.90±2.59***
<i>M.alba</i> (6)	200	142.4±1.81***
<i>M.alba</i> (6)	400	124.40±1.64***
Imipramine (6)	10	108.00±2.07***

Values are given as mean ±SEM (N=6 in each group), ***P<0.001, as compared to control





Table 2: Effect of *Morus alba* leaves extract on immobility time in Forced swimming test

Treatment	DOSE (mg/kg)	Immobility time (in sec)
Vehicle Control(6)	-	145.40±3.84
<i>M.alba</i> (6)	100	133.00±2.12***
<i>M.alba</i> (6)	200	124.20±1.42***
<i>M.alba</i> (6)	400	112.60±1.94***
Imipramine (6)	10	91.20±2.32***

Values are given as mean ±SEM (N=6 in each group), ***P<0.001, as compared to control



Ample amount of free radicals are constantly produced in the body through different biochemical reactions, if the level of these free radicals surpasses the normal value, it leads to oxidative impairment in cells as well as tissues leading to various degenerative diseases in the body. Plant extracts are frequently used as natural source of antioxidants which can prevent these degenerative disorders. Antioxidants neutralise the free radicals in the body which improves the disease conditions. Chlorogenic and gallic acid are antioxidants present in the plant *Morus alba*, which is demonstrated in the article by using phytochemical screening and HPTLC fingerprinting. These in-vitro antioxidants can fight against free radicals produced in the body and hence results in improvements in disease conditions of depression and other ailments like diabetes and cardiovascular diseases. Through our results it was demonstrated that *morus alba* leaves are may be acceptable as a natural antioxidant and antidepressant with various bioactive components used for the treatment of numerous other ailments.

REFERENCES:

1. Thaipitakwong, T.; Numhom, S.; Aramwit, P. Mulberry leaves and their potential effects against cardiometabolic risks: a review of chemical compositions, biological properties and clinical efficacy. *Pharmaceutical Biology*; 2018; 56(1); 109-118.
2. Sharma, A.; Krishna, V.; Kaur, P.; Rayal, R. Characterization and screening of various Mulberry varieties through morpho-biochemical characteristics. *Journal of Global Biosciences*; 2015; 4(1); 1186-1192.
3. Killedar, S. G.; Pawar, A. V. Preparation of herbal tea from mulberry leaves. *Journal of Medicinal Plants Studies*; 2017; 5(2): 325-328.
4. Kim, S. B.; Chang, B. Y.; Hwang, B. Y.; Kim, S. Y.; Lee, M. K. Pyrrole alkaloids from the fruits of *Morus alba*. *Bioorganic & Medicinal Chemistry Letters*; 2014; 24(24), 5656-5659.
5. Asano, N.; Yamashita, T.; Yasuda, K.; Ikeda, K.; Kizu, H.; Kameda, Y.; Ryu, K. S. Polyhydroxylated Alkaloids Isolated from Mulberry Trees (*Morus alba*L.) and Silkworms (*Bombyx mori*L.). *Journal of Agricultural and Food Chemistry*, 2001; 49(9), 4208-4213.
6. Wang, X.; Kang, J.; Wang, H-Q.; Liu, C.; Li, B-M.; Chen, R-Y. Three new alkaloids from the fruits of *Morus alba*, *Journal of Asian Natural Products Research*, 2014; 16(5), 453-458.
7. Chu, Q.; Lin, M.; Tian, X.; Ye, J. Study on capillary electrophoresis-amperometric detection profiles of different parts of *Morus alba* L. *J. Chromatogr. A* 2006, 1116, 286-290.
8. De Freitas, M. M.; Fontes, P. R.; Souza, P. M.; William Fagg, C.; Neves Silva Guerra, E.; de Medeiros Nóbrega, Y. K.; Silveira, D.; Fonseca-Bazzo, Y.; Alberto Simeoni, L.; Homem-de-Mello, M.; Oliveira Magalhães, P. Extracts of *Morus nigra* L. leaves standardized in chlorogenic acid, rutin and isoquercitrin: tyrosinase inhibition and cytotoxicity. *Plos One*, 2016; 11(9), e0163130.
9. Kokate, C.K.; *Practical Pharmacognosy*. Delhi: VallabhPrakashan; 2008.p. 108-9.
10. AOAC. *Official Methods of Analysis* .15th ed. Washington, VA: Association of Official Analytical Chemists; 1990.
11. Doshi, G. M.; Zine, S. P.; Chaskar, P. K.; Une, H. D. Solicitation of HPLC and HPTLC Techniques for Determination of Rutin from *Polyalthialongifolia Thwaites*, *harmacognosy Res.* 2014; 6(3): 234-239.
12. Chermat, R.; Thierry, B.; Micro, J.A.; Steru, L.; Simon, P. Adaptation of the tail suspension test of the rat. *J Pharmacol* 1986; 17:348-350.
13. Takamori, Tadano, T.; Yoshida, S.; Okuyama, S. Repeated treatment with imipramine, fluvoxamine and tranlycypromine decreases the number of escape failures by activating dopaminergic systems in a rat learned helplessness test *Life Sci*; 2001; 69:1919-1926.