

# Effect of Carbamazepine on Reproductive Function in Female Wistar Rats

Oyededeji K.O.\*<sup>1</sup>, Umar B.F.<sup>1</sup>

<sup>1</sup>Department of Physiology, College of Medicine and Health Sciences,  
Afe Babalola University, Ado-Ekiti, Nigeria.

\* sinaoyededeji@yahoo.com

## Abstract

**Introduction:** This study was designed to investigate the effect of carbamazepine on reproductive function in female Wistar rats.

**Methods:** Eighteen female rats (120 – 160 g) were used for the estrous cycle and histopathological studies. Carbamazepine (5.71 mg/kg) was administered orally on daily basis for 21 and 50 days respectively for the estrous cycle and histological studies. Estrous cycle was carried out using the technique of Marcondes *et al.*, histologies of the ovaries and uteri were also carried out. Data were analysed using descriptive statistics and student's t-test at  $p=0.05$ .

**Results:** Treatment of rats for 21 days with carbamazepine (5.71 mg/kg) produced a significant ( $p<0.05$ ) decrease in proestrous phase of the estrous cycle relative to the respective control. The histopathological study revealed that treatment of rats with carbamazepine for 50 days produced no visible lesion on the ovaries and uteri, which is similar to what was observed in the control rats.

**Conclusion:** It can therefore be concluded that carbamazepine probably has pro-fertility effect, and probably induced no deleterious effect on the ovaries and uteri in female Wistar rats.

**Keywords:** Carbamazepine, Proestrous, Diestrous, Endometrium, Rats.

## INTRODUCTION

Carbamazepine is an anticonvulsant medication used primarily in the treatment of epilepsy and neuropathic pain [1]. It is used in schizophrenia along with other medications and as a second-line agent in bipolar disorder [2]. Carbamazepine appears to work as well as phenytoin and valproate for focal and generalized seizures [3].

The anticonvulsant effect of carbamazepine on spontaneous seizures in rats with kainate-induced epilepsy has been reported [4]. The teratogenic effects of carbamazepine administration in pregnant rats [5] and mice [6] have been reported. The harmful effect of carbamazepine on the postnatal development of the rat ventral prostate has been reported [7]. Analgesic synergism of gabapentin and carbamazepine in rat model of diabetic neuropathic pain has been reported [8]. The neurobehavioural and cognitive changes induced by carbamazepine and/or phenytoin in Wistar rats have been reported [9]. Carbamazepine induced changes in maternal reproductive performance and fetal growth retardation in rats have also been reported [10].

However, due to scanty information from literature on the effect of carbamazepine on reproductive parameters in female rats, this study therefore aims at investigating the effect of this anticonvulsant agent on these aforementioned parameters in female rats.

## MATERIALS AND METHODS

### Experimental Animals

Adult female rats weighing between 120 g – 160 g bred in the Pre-Clinical Animal House of the College of Medicine and Health Sciences, Afe Babalola University were used. They were housed under standard laboratory conditions and had free access to feed and water; they were acclimatized for two weeks to laboratory conditions before

the commencement of the experiments. All experiments were carried out in compliance with the recommendations of Afe Babalola University Ethics Committee on guiding principles on care and use of animals.

### Drug

Carbamazepine (Hovid Pharmacy, Ltd) was bought from Danax Pharmacy, Ibadan, Nigeria.

Carbamazepine (200 mg) was dissolved in 10 ml of distilled water to give a concentration of 20 mg/ml.

The dosage of carbamazepine used in this study was in accordance with that reported by the manufacturer.

### Experimental Design

#### Study on Estrous Cycle

Six matured female rats showing at least three regular 4 – 5 day cycles were used for this study. Vaginal lavages (smears) were examined microscopically everyday at a constant interval of 4.30 – 5.30 p.m. for 21 days before and after treatments with the anticonvulsant agent. The smears were classified into one of the phases of the estrous cycle using the Marcondes technique [11]. Vaginal secretion was collected with a plastic pipette filled with 10  $\mu$ L of normal saline (NaCl 0.9 %) by inserting the tip into the rat's vagina, but not deeply. Vaginal fluid was placed on glass slide. One drop was collected with a clean tip from each rat. Unstained material was observed under a light microscope, without the use of condenser lens, with x10 and x40 objective lenses. Three types of cells could be recognized: round and nucleated ones are epithelial cells; irregular ones without nucleus are the cornified cells; and the little round ones are the leucocytes. The proportion (preponderance) among them was used for the determination of estrous cycle phases [12, 13]. The duration of the estrous cycle was determined. In this study,

the experimental animals also served as the control. The first 21 days served as the control days, while the last 21 days served as the treatment days. Each of the 6 rats for this estrous cycle study received 5.71 mg/kg of carbamazepine (orally).

### Histopathological Study

In another set of experiment, twelve matured female rats divided into two equal groups (six animals per group) received the following treatments of the anticonvulsant agent and control (orally) per day for fifty days as follows: Group I rats received 0.5 ml/100 g of distilled water as the control group.

Group II rats received 5.71 mg/kg of carbamazepine.

On the 51st day, all the rats were sacrificed by an overdose of chloroform. The ovaries and uteri were dissected out, cleaned of fat and immediately fixed in Bouin's fluid.

### Histological preparation of tissues

After weighing the ovaries and uteri, they were immediately fixed in Bouin's fluid for 12 hours and the Bouin's fixative was washed from the samples with 70 % alcohol. The tissues were then cut in slabs of about 0.5 cm transversely and the tissues were dehydrated by passing through different grades of alcohol: 70 % alcohol for 2 hours, 100 % alcohol for 2 hours, and finally 100 % alcohol for 2 hours. The tissues were then cleared to remove the alcohol, the clearing was done for 6 hours using xylene. The tissues were then infiltrated in molten paraffin wax for 2 hours in an oven at 57°C, thereafter the tissues were embedded. Serial sections were cut using rotary microtome at 5 microns (5 µm). The satisfactory ribbons were picked up from a water bath (50 - 55°C) with microscope slides that had been coated on one slide with egg albumin as an adhesive and the slides were dried in an oven. Each section was deparaffinized in xylene for 1 minute before immersed in absolute alcohol for 1 minute and later in descending grades of alcohols for about 30 seconds each to hydrate it. The slides were then rinsed in water and immersed in alcoholic solutions of hematoxylin for about 18 minutes. The slides were rinsed in water, and then differentiated in 1 % acid alcohol and then put inside a running tap water to blue and then counterstained in alcoholic eosin for 30 seconds and rinsed in water for a few seconds, before being immersed in 70 %, 90 % and twice in absolute alcohol for 30 seconds each to dehydrate the preparations. The preparations were cleared of alcohol by dripping them in xylene for 1 minute. Each slide was then cleaned, blotted and mounted with DPX and cover slip, and examined under the microscope. Photomicrographs were taken at x40 and x100 magnifications.

### Statistical Analysis


The mean and standard error of mean (S.E.M.) were calculated for all values. Comparison between the control and the treated group was done using student's t-test. Differences were considered statistically significant at  $p < 0.05$ .

## RESULTS


Treatment of rats for 21 days with carbamazepine (5.71 mg/kg) produced a significant ( $p < 0.05$ ) decrease in proestrous phase, but caused insignificant ( $p > 0.05$ ) changes in the estrous, metestrous and diestrous phases of the estrous cycle relative to their respective controls (Figure 1).

Treatment of rats with carbamazepine (5.71 mg/kg) for 50 days produced no visible lesion on the ovaries, which is similar to what was observed in the control rats (Plates 1 and 2).

Treatment of rats with carbamazepine (5.71 mg/kg) for 50 days produced no pathological effect on the uteri, which is similar to what was observed in the control rats (Plates 3 and 4).




**Figure 1: Effect of 21 days treatment with carbamazepine on estrous cycle (n = 6, \* $p < 0.05$ )**




**Plate 1: Effect of 0.5 ml/100 g distilled water (control) on the ovary at x100.**


Photomicrograph showing a normal ovary (O) with no visible lesion seen.



**Plate 2:** Effect of carbamazepine (5.71 mg/kg) on the ovary at x100.  
Photomicrograph showing an ovary with no pathological lesion (N).



**Plate 3:** Effect of 0.5 ml/100 g distilled water (control) on the uterus at x100.  
Photomicrograph showing normal endometria (E) and myometrium (M) with no visible lesion seen.



**Plate 4:** Effect of carbamazepine (5.71 mg/kg) on the uterus at x100  
Photomicrograph showing normal endometrial (E) and myometrial (M) with no pathological lesion present.

## DISCUSSION

The estrous cycle study revealed that carbamazepine caused a significant change in the duration of a phase of the estrous cycle. Contrary report was given by [14] in *Portulaca oleracea* extracts treated rats. This suggests that the anticonvulsant agent caused imbalances of the ovarian and extraovarian hormones, since it has been reported that imbalance in these hormones lead to irregularity in the ovarian functions and duration of the estrous cycle [15].

Treatment of rats with carbamazepine caused a significant decrease in proestrous phase of the estrous cycle which suggests the maturation of the follicles in the pre-ovulatory phase was hastened, *vis-à-vis*, leading to maturation of the Graafian follicles. Contrary result was reported by [16] in alcohol treated rats.

The ovarian photomicrographs of the carbamazepine treated rats presented with no pathologic lesion which suggests the non-toxic effect of the drug on the ovaries at histologic level. Similar result was reported by [14] in *Portulaca oleracea* treated rats.

The uterine photomicrographs of the carbamazepine treated rats presented with no pathologic lesion which probably indicates the non-toxic effect of the drug on the uteri at histologic level. Similar result was reported by [17] in *Allium sativum* extract treated rats.

In conclusion, this study has shown that carbamazepine probably has pro-fertility effect in female Wistar rats. It also revealed that carbamazepine probably induced no deleterious effect on the ovaries and uteri at histological level level in female Wistar rats. However, the effect of this anticonvulsant agent on human reproductive function is unknown; nevertheless, considering these findings in animal model, it is recommended that women with infertility problems could take carbamazepine for infertility therapeutic purpose.

## Conflict of Interest

We vehemently declare that there is no conflict of interests in this research work.

## REFERENCES

- [1] The American Society of Health-System Pharmacists. 'Carbamazepine'. Archived from the original on 2015-02-27. Retrieved 28 March, 2015.
- [2] Nevitt SJ, Marson AG, Weston J, Tudur SC. Sodium valproate versus phenytoin monotherapy for epilepsy: an individual participant data review. The Cochrane Database of System Rev. 2018; 8: CD001769.
- [3] Nevitt SJ, Marson AG, Tudur SC. Carbamazepine versus phenytoin monotherapy for epilepsy: an individual participant data review. The Cochrane Database of System Revs. 2019; 7: CD001911.
- [4] Grabenstatter HL, Clark S, Dudek FE. Anticonvulsant effect of carbamazepine on spontaneous seizures in rats with kainate-induced epilepsy: comparison of intraperitoneal injections with drug-in-food protocols. *Epilepsia*. 2007; 48 (12): 2287–2295.
- [5] El-Gaafarawi I, Abouel-Magd M. Teratogenic effect of carbamazepine administration in pregnant rats. *The Egyptian J Hospit Med*. 2015; 59: 244- 257.
- [6] Jose M, Sreelatha HV, James MV, Arumughan S, Thomas SV. Teratogenic effects of carbamazepine in mice. *Annals of Indian Acad Neuro*. 2017; 20 (2): 132-137.
- [7] Oliva SU, Scarano WR, Okada FK, Miraglia SM. Harmful effect of carbamazepine on the postnatal development of the rat ventral prostate. *Reprod Biol Endo*. 2012; 10: 22 (2012).

- [8] AL-Mahmood SMA, Abdullah STBC, Ahmad NNFN, Mohamed AHB, Abdul Razak T. Analgesic synergism of gabapentin and carbamazepine in rat model of diabetic neuropathic pain. *Trop J Pharm Res.* 2016; 15 (6): 1191-1195.
- [9] Aliyu H, Ayo JO, Ambali SF, Zezi AU, Kobo PI, Uchendu C. Evaluation of neurobehavioural and cognitive changes induced by carbamazepine and/or phenytoin in Wistar rats. *Global J Med Res: A Neurol and Nervous Syst.* 2016; 16 (1) Version 1.0.
- [10] Nunes M, Coelho de Sousa FD, Andretta RR, Miraglia SM, Urban de Oliva S. Carbamazepine causes changes in maternal reproductive performance and fetal growth retardation in rats. *bioRxiv.* doi: <https://doi.org/10.1101/2020.09.18.30348>.
- [11] Marcondes FK, Bianchi FJ, Tanno AP. Determination of the estrous cycle phases of rats: some helpful considerations. *Braz J Bio.* 2002; 62 (4a): 609 – 614.
- [12] Long JA, Evans HM. The estrous cycle in the rat and its associated phenomena. *Memo Uni Califor.* 1922; 6: 1-148.
- [13] Mandl AM. The phases of the oestrous cycle in the adult white rat. *J Exptal Bio.* 1951; 28: 576-5 84.
- [14] Oyedemi KO, Bolarinwa AF. Effects of extracts of *Portulaca oleracea* on reproductive functions in female albino rats. *Afr J Biomed Res.* 2010; 13: 213-218.
- [15] Circosta C, Sanogo R, Occhiuto F. Effects of *Calotropis procera* on estrous cycle and on estrogenic functionality in rats. *Farmacol.* 2001; 56: 373-378.
- [16] Oyedemi KO, Bolarinwa AF, Fashina AM. Effect of alcohol consumption on hematological and reproductive parameters in female albino rats. *IOSR J Dent Med Sci.* 2013; 3 (5):76-79.
- [17] Raji LO, Fayemi OE, Ameen SA, Jagun AT. The effects of aqueous extract of *Allium sativum* (Garlic) on some aspects of reproduction in the female albino rats. *Global Vet.* 2012; 8 (4): 414-420.